
Contrast 5 To McGray 1st Company Inc. Permanon required by regordation to display



Periods of Western Astronomy

- Western astronomy divides into 4 periods
 - Prehistoric (before 500 B.C.)
 - Cyclical motions of Sun, Moon and stars observed
 - Keeping time and determining directions develops
 - Classical (500 B.C. to A.D. 1400)
 - Measurements of the heavens
 - Geometry and models to explain motions
 - Renaissance (1400 to 1650)
 - Accumulation of data led to better models
 - Technology (the telescope) enters picture
 - Modern (1650 to present)
 - Physical laws and mathematical techniques
 - Technological advances accelerate


Ancient Greek Astronomers

Moon Earth Venus Sun Mars Jupiter

- Through the use of models and observations, they were the first to use a careful and systematic manner to explain the workings of the heavens
- Limited to naked-eye observations, their idea of using logic and mathematics as tools for investigating nature is still with us today
- Their investigative methodology is in many ways as important as the discoveries themselves

Early Ideas: Pythagoras

• Pythagoras taught as early as 500 B.C. that the Earth was round, based on the belief that the sphere is the perfect shape used by the gods



Early Ideas: Aristotle

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display




Copyright 2009 Anthony Ayiomamitis

- By 300 B.C., Aristotle presented naked-eye observations for the Earth's spherical shape:
 - Shape of Earth's shadow on the Moon during an eclipse

Early Ideas: Aristotle


Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



He also noted that a traveler moving south will see stars previously hidden by the southern horizon


Early Ideas: The Size of the Earth

- Eratosthenes (276-195 B.C.) made the first measurement of the Earth's size
- He obtained a value of 25,000 miles for the circumference, a value very close to today's value




Early Ideas: The Size of the Earth

 He measured the shadow length of a stick set vertically in the ground in the town of Alexandria on the summer solstice at noon, converting the shadow length to an angle of solar light incidence, and using the distance to Syene, a town where no shadow is cast at noon on the summer solstice




Early Ideas: Distance and Size of the Sun and Moon


- The sizes and distances of the Sun and Moon relative to Earth were determined by Aristarchus about 75 years before Eratosthenes measured the Earth's size
- Once the actual size of the Earth was determined, the <u>absolute</u> sizes and distances of the Sun and Moon could be determined



Early Ideas: Distance and Size of the Sun and Moon

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.






• These <u>relative</u> sizes were based on the *angular size* of objects and a simple geometry formula relating the object's diameter, its angular size, and its distance

Early Ideas: Distance and Size of the Sun and Moon

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



• Aristarchus, realizing the Sun was very large, proposed the Sun as center of the Solar System, but the lack of *parallax* argued against such a model

Measuring the Diameter of Astronomical Objects


Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display

$$l = 2\pi d \times \frac{\alpha}{360^{\circ}}$$

l – linear size of object


d – distance to object

 α – angular size of object



Planets and the Zodiac

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




- The planets (Greek for "wanderers") do not follow the same cyclic behavior of the stars
- The planets move relative to the stars in a very narrow band centered about the ecliptic and called the *zodiac*
- Motion and location of the planets in the sky is a combination of all the planets' orbits being nearly in the same plane and their relative speeds about the Sun

Planets and the Zodiac

Apparent motion
 of planets is
 usually from west
 to east <u>relative to</u>
 the stars, although
 on a daily basis,
 the planets always
 rise in the east

Retrograde Motion






- Occasionally, a planet will move from east to west relative to the stars; this is called *retrograde motion*
- Explaining retrograde motion was one of the main reasons astronomers ultimately rejected the idea of the Earth being located at the center of the solar system

Early Ideas: The Geocentric Model


- Because of the general east to west motion of objects in the sky, *geocentric theories* were developed to explain the motions
- Eudoxus (400-347 B.C.) proposed a geocentric model in which each celestial object was mounted on its own revolving transparent sphere with its own separate tilt
- The faster an object moved in the sky, the smaller was its corresponding sphere
- This simple geocentric model could not explain retrograde motion without appealing to clumsy and unappealing contrivances

Early Ideas: The Geocentric Model




Ptolemy of Alexandria

- Ptolemy of Alexandria improved the geocentric model by assuming each planet moved on a small circle, which in turn had its center move on a much larger circle centered on the Earth
- The small circles were called *epicycles* and were incorporated so as to explain retrograde motion



Ptolemy of Alexandria

- Ptolemy's model was able to predict planetary motion with fair precision
- Discrepancies remained and this led to the development of very complex Ptolemaic models up until about the 1500s
- Ultimately, all the geocentric models collapsed under the weight of "Occam's razor" and the *heliocentric models* prevailed




Non-Western Contributions


- Islamic Contributions
 - Relied on celestial phenomena to set its religious calendar
 - Created a large vocabulary still evident today (e.g., zenith,
 Betelgeuse)
 - Developed algebra and Arabic numerals
- Asian Contributions
 - Devised constellations based on Asian mythologies
 - Kept detailed records of unusual celestial events (e.g., eclipses, comets, supernova, and sunspots)
 - Eclipse predictions

- Nicolaus Copernicus (1473-1543)
 - Could not reconcile centuries of data with Ptolemy's geocentric model
 - Consequently, Copernicus reconsidered Aristarchus's heliocentric model with the Sun at the center of the solar system


Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



- Heliocentric models
 explain retrograde motion
 as a natural consequence
 of two planets (one being
 the Earth) passing each
 other
- Copernicus could also derive the relative distances of the planets from the Sun



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



- However, problems remained:
 - Could not predict planet positions any more accurately than the model of Ptolemy
 - Could not explain lack of parallax motion of stars
 - Conflicted withAristotelian "common sense"

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display



- Tycho Brahe (1546-1601)
 - Designed and built instruments of far greater accuracy than any yet devised
 - Made meticulous measurements of the planets

- Tycho Brahe (1546-1601)
 - Made observations

 (supernova and comet) that
 suggested that the heavens
 were both changeable and
 more complex than
 previously believed
 - Proposed compromise geocentric model, as he observed no parallax motion!

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




- Johannes Kepler (1571-1630)
 - Upon Tycho's death,his data passed toKepler, his youngassistant
 - Using the very precise
 Mars data, Kepler
 showed the orbit to be
 an *ellipse*




Johannes Kepler (1571-1630)

- Using Tycho Brahe's data, discovered that planets do not move in circles around the Sun, rather, they follow ellipses with the Sun located at one of the two foci!
- Astronomers use the term *eccentricity* to describe how round or "stretched out" an ellipse is the higher (closer to 1) the eccentricity, the flatter the ellipse.









Kepler's 1st Law

• Planets move in elliptical orbits with the Sun at one *focus* of the ellipse



Kepler's 2nd Law

- The orbital speed of a planet varies so that a line joining the Sun and the planet will sweep out equal areas in equal time intervals
- The closer a planet is to the Sun, the faster it moves




Kepler's 3rd Law

- The amount of time a planet takes to orbit the Sun is related to its orbit's size
- The square of the period, P, is proportional to the cube of the *semimajor* axis, a

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

$$P^2$$
 years = a^3 AU



P = time to complete orbita = semimajor axis


C

Kepler's 3rd Law

- This law implies that a planet with a larger average distance from the Sun, which is the semimajor axis distance, will take longer to circle the Sun
- Third law hints at the nature of the force holding the planets in orbit

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

$$P^2$$
 years = a^3 AU



P = time to complete orbita = semimajor axis


C

Kepler's 3rd Law

• Third law can be used to determine the semimajor axis, a, if the period, P, is known, a measurement that is not difficult to make

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


$$P^2$$
 years = a^3 AU




P = time to complete orbita = semimajor axis

C


- Galileo (1564-1642)
 - Contemporary of Kepler
 - First person to use the telescope to study the heavens and offer interpretations
 - The Moon's surface has features similar to that of the Earth ⇒ The Moon is a ball of rock



- The Sun has spots ⇒ The Sun is not perfect, changes its appearance, and rotates
- Jupiter has four objects
 orbiting it ⇒ The objects are
 moons and they are not
 circling Earth
- Milky Way is populated by uncountable number of stars
 ⇒ Earth-centered universe is too simple




Evidence for the Heliocentric Model




• Venus undergoes full phase cycle ⇒ Venus must circle Sun

- Credited with originating the experimental method for studying scientific problems
- Deduced the first correct "laws of motion"
- Was brought before the Inquisition and put under house arrest for the remainder of his life




Isaac Newton

- Isaac Newton (1642-1727) was born the year Galileo died
- He made major advances in mathematics, physics, and astronomy



Isaac Newton

- He pioneered the modern studies of motion, optics, and gravity and discovered the mathematical methods of calculus
- It was not until the 20th century that Newton's laws of motion and gravity were modified by the theories of relativity



The Growth of Astrophysics

New Discoveries

- In 1781, Sir William Herschel discovered Uranus;
 he also discovered that stars can have companions
- Irregularities in Uranus's orbit together with law of gravity led to discovery of Neptune

New Technologies

- Improved optics led to bigger telescopes and the discovery of nebulas and galaxies
- Photography allowed the detection of very faint objects